

Document ID:	TDS-MCP-070-50L	Version:	001
Date of Issue:	10-JUL-2022	Approved by:	Dr. Iman Kamranfar
Review Date:	10-JAN-2024	Signature:	Har
Title:	TECHNICAL DATA SHEET		

-Glutamine, without Sodium onate and Phenol Red 170-50L 4 years 2 to 8 °C
170-50L 4 years
4 years
2 to 8 °C
2100 C
ambient
Concentration (mg/L)
90.00
60.50
20.00
30.00
58.00
26.09
50.00
292.00
20.30
50.00
75.00
87.50
15.00
25.00
30.00
10.00
40.00
10.00
35.00
50.00
0.50
1.00
1.50
1.00
0.01
1.00
2.00
0.01
0.01 2.00
0.01 2.00 1.00
0.01 2.00 1.00 1.00
0.01 2.00 1.00 1.00 0.10
0.01 2.00 1.00 1.00 0.10 1.00
0.01 2.00 1.00 1.00 0.10



Document ID:	TDS-MCP-070-50L	Version:	001
Date of Issue:	10-JUL-2022	Approved by:	Dr. Iman Kamranfar
Review Date:	10-JAN-2024	Signature:	And
Title:	TECHNICAL DATA SHEET		

Other Components	
Dextrose (D-Glucose)	2000.00
Reduced Glutathione	0.05
Methyl Linoleate	0.03
Sodium Pyruvate	25.00
Inorganic Salts	
Calcium Chloride 2H ₂ O	264.90
Cupric Sulfate	0.00006
Ferric Nitrate Nonahydrate [Fe(NO ₃) ₃ 9H ₂ O]	0.0001
Magnesium Sulfate, Anhydrous	97.70
Potassium Chloride	400.00
Sodium Chloride	6800.00
Sodium Phosphate Monobasic Monohydrate (NaH ₂ PO ₄ H ₂ O)	140.00
Zinc Sulfate Heptahydrate (ZnSO ₄ 7H ₂ O)	0.0002
Gram of Powder required for 1 L Medium	10.814

Specifications		
Appearance	Off-white powder	
Solubility	10.814 g/L in ≤ 30 minutes	
Moisture Content	≤ 2.0 %	
pH without NaHCO3	Test and report	
pH with NaHCO3	6.8 - 7.8	
Osmolality without NaHCO3	240-280 mM/kg	
Osmolality with NaHCO3 (2.2 g/L)	280-340 mM/kg	
Endotoxin	≤ 1.0 EU/ml	
Mycoplasma	Not detected	
	,	

Note: Sodium Bicarbonate is not included in the powdered media (add 2200.00 mg/L).

Product description

Willliams et al., (1971)¹ introduced a new method based on sequential plating technique to more efficiently isolate and culture new born epithelial liver cells. They used a modified version of MEM medium called Williams' Medium D which was enriched in amino acids and double glucose content. Williams and Gunn (1974)² conducted further studies led to release of Williams' Medium E to be used for the effective long-term culture of adult liver cells. During last decades Williams' Medium E has been widely used for the culture of liver epithelial cells, as well as primary hepatocytes from different species (e.g., human HepaRG cells).

William's Medium E contains unique ingredients such as zinc, iron, manganese, non-essential amino acids, the reducing agent glutathione and the lipid methyl linoleate. William's E Medium can be supplemented, usually with 5-10% fetal bovine serum may be required. William's E Medium uses a sodium bicarbonate (2.2 g/l) buffer system and therefore requires a 5-10% CO 2 environment to maintain physiological pH.

References

- 1. Williams, G.M., and Gunn, J.M., Long-Term Cell Culture of Adult Rat Liver Epithelial Cells. Exp. Cell Research, 89, 139-142 (1974).
- 2. Williams, G.M. et al., Isolation and Long-Term Cell Culture of Epithelial-Like Cells From Rat Liver. Exp. Cell Research, 69, 106-112 (1971).

THIS PRODUCT IS FOR LABORATORY USE ONLY.