

Document ID:	TDS-S-FBS-CO-055	Version:	001
Date of Issue:	10-JAN-2023	Approved by:	Dr. Iman Kamranfar
Review Date:	10-JAN-2025	Signature:	H
Title:	TECHNICAL DATASHEET		

# **Foetal Bovine Serum**

Filtration, Treatment	Ultra low IgG
Origin	Colombia
Product Code	S-FBS-CO-055
Shelf Life	5 years from DOM
Storage Temperature	<-15°C
Shipping Temperature	dry ice

# **QC Specifications**

Physical and Chemical Analysis	Method	Specifications	Units
Appearance	Visual	Clear yellow-amber	n/a
Identity	Internally Validated	Bovine	n/a
pH at RT	Electronic pH Meter	6.8 - 8.2	n/a
Osmolality	Osmometer	260 - 340	mOsm/kg
Endotoxin	LAL Kinetic	< 10.0	EU/ml
Free Hemoglobin	Colorimetric	< 25.0	mg/dl
Specific Gravity	Mass Balance	> 1.01	g/ml
Sterility			
Sterility	Internally Validated	Pass	n/a
Mycoplasma	gPCR	Not detected	n/a
тусоргазита	qr civ	Not detected	11/4
Protein Profile			
Total Protein	IDEXX Catalyst One	3.0 - 4.5	g/dl
Albumin	IDEXX Catalyst One	1.4 - 3.4	g/dl
Globulin	IDEXX Catalyst One	0.4 - 2.4	g/dl
IgG	ELISA	<5	μg/ml
Electrophoretic Pattern	Capillary Electrophoresis	Normal	n/a
- F	, , , , , , , , , , , , , , , , , , ,	<u> </u>	1 .4
Antibiotic Testing			
Tetracycline	IDEXX Snap Test	Test and report	n/a
Oxytetracycline	IDEXX Snap Test	Test and report	n/a
Chlortetracycline	IDEXX Snap Test	Test and report	n/a
Virus Testing	•		
BVDV/BHV-1/PI-3 (CPE)	Cell Culture	Not detected	n/a
Rabies Virus	qPCR	Not detected	n/a
Bluetongue Virus (BTV)	qPCR	Not detected	n/a
BRSV	qPCR	Not detected	n/a
Reo Virus	qPCR	Not detected	n/a
BAV	qPCR	Not detected	n/a
BoPV-1, -2	qPCR	Not detected	n/a
Antibody Testing			1
BVDV - Antibody Type 1	Serum Neutralization Test (Cell Culture) or Detection of Antibodies (ELISA)	Test and report	n/a
BVDV - Antibody Type 2	Serum Neutralization Test (Cell Culture) or Detection of Antibodies (ELISA)	Test and report	n/a
BHV-1	Detection of Antibodies (ELISA)	Test and report	n/a
PI-3	Detection of Antibodies (ELISA)	Test and report	n/a
Biochemistry			
Aspartate Aminotransferase (AST)	IDEXX Catalyst One	Record	U/L
Alanine Aminotransferase (ALT)	IDEXX Catalyst One	Record	U/L
, warmie , willing a distribute (ALI)	IDEAN Cutulyst Offic	necoru	0/ L



Document ID:	TDS-S-FBS-CO-055	Version:	001
Date of Issue:	10-JAN-2023	Approved by:	Dr. Iman Kamranfar
Review Date:	10-JAN-2025	Signature:	H
Title:	TECHNICAL DATASHEET		

Alkaline Phosphatase (ALKP)	IDEXX Catalyst One	Record	U/L
Gamma-Glutamyl Trans.(GGT)	IDEXX Catalyst One	Record	U/L
, , ,	•	Record	mmol/L
Cholesterol (CHOL)	IDEXX Catalyst One		mmol/L
Glucose (GLU)	IDEXX Catalyst One	Record	· '
Urea (BUN)	IDEXX Catalyst One	Record	mmol/L
Creatinine (CREA)	IDEXX Catalyst One	Record	μmol/L
Uric Acid (URIC)	IDEXX Catalyst One	Record	μmol/L
Calcium (Ca)	IDEXX Catalyst One	Record	mmol/L
Phosphorus (PHOS)	IDEXX Catalyst One	Record	mmol/L
Total Bilirubin (TBIL)	IDEXX Catalyst One	Record	μmol/L
Magnesium (Mg)	IDEXX Catalyst One	Record	mmol/L
Sodium (Na)	IDEXX Catalyst One	Record	mmol/L
Potassium (K)	IDEXX Catalyst One	Record	mmol/L
Chloride (CI)	IDEXX Catalyst One	Record	mmol/L
Cell Culture Testing - Option 1			
Cell Line	Method	Specifications	Results
L-929, HeLa, MRC-5	Morphology	Tested vs. Control Serum	Scoring
2 323, 11624, 14116 3	Weighteregy	rested vs. control serum	System 1
L-929, HeLa, MRC-5	Density	Tested vs. Control Serum	Scoring System 2
L-929, HeLa, MRC-5	Cell Count	Cell count [log10/ml]/dead cells vs. Control	Record
Scoring system	Meaning		Results
<u> </u>	Dead Cells		0
	Many Cells degenerate		
	and many dead cells		1
	Cells partially		
4. Marris India	degenerate and many		2
1 - Morphology	dead cells		
	Few cells degenerate		2
	and few dead cells		3
	Without pathological		4
	findings		4
	Single cells/cell		
	aggregates		0
	< 50% confluency		1
2 - Density	50 - 90% confluency		2
	confluency		3
	overly confluent		4
		•	
Cell Culture Testing - Option 2			
Cell Line	Method	Specifications	Units
	Multiple Passage -		
BUW 24 - 120 5	Records results	- 750/ of a sales land	0/
BHK-21, MRC-5	vs. control at day: 0, 3,	>75% of control growth	%
	6, 12		
	Plating Efficiency -		
BUW 24 . 120 5	Records results	. 750/ P5	0/
BHK-21, MRC-5	vs. control at day: 0, 3,	>75% PE vs. control PE	%
	6, 12		
	Cloning Efficiency -		
DUIK 24 MADG 5	Records results	. 750/ 65 65	0/
BHK-21, MRC-5	vs. control at day: 0, 3,	>75% CE vs. control CE	%
	6, 12		

### **RECOMMENDED USE:**

## Storage

To effectively preserve the integrity of animal serum, it should be stored frozen and protected from light. The recommended storage temperature is <-15°C.



Document ID:	TDS-S-FBS-CO-055	Version:	001
Date of Issue:	10-JAN-2023	Approved by:	Dr. Iman Kamranfar
Review Date:	10-JAN-2025	Signature:	Mar
Title:	TECHNICAL DATASHEET		

Multiple thaw/freeze cycles should be avoided, as they will accelerate the degradation of serum nutrients and can encourage the formation of insoluble precipitates. For this reason, serum should never be stored in "frost-free" freezers. These types of freezers periodically warm themselves to avoid internal ice deposits and are detrimental to the stability of frozen serum products.

### **Suggested Thawing Procedure**

- 1. Remove the serum bottles from the freezer and allow them to adjust to room temperature for approximately 10 minutes.
- 2. Place each container in a 30 to 37 °C water bath or incubator. Excessive temperatures will degrade heat labile nutrients. If using a water bath, prevent the bottle caps from being submerged.
- 3. Gently agitate the bottles every 10 15 minutes until the serum is completely thawed.

#### **Efficient and Effective Usage**

After thawing, use the serum promptly. Liquid serum may be stored refrigerated (2 to 8 °C) up to four weeks. To avoid thaw/freeze cycles or long periods of refrigeration, it is recommended that any unused serum be immediately dispensed into small, useful aliquots and refrozen for future use.

Product use: This product is not intended for human or animal consumption or therapeutic use.